

Bounds on Broken R -Parity from Leptonic Meson Decays

Herbi Dreiner^{1*}, Giacomo Polesello^{2†}, Marc Thormeier^{3‡}

¹ *Physikalisches Institut der Universität Bonn,
Nußallee 12, 53115 Bonn, Germany*

² *INFN, Sezione di Pavia,
Via Bassi 6, 27100 Pavia, Italy*

³ *Department of Theoretical Physics, University of Oxford,
1 Keble Road, Oxford OX1 3NP, United Kingdom*

Abstract

Investigating leptonic decays of π^- , K^- , B^- , π^0 , K_L^0 , B_s^0 we present new bounds on some products of two R -parity violating coupling constants. For mesons of a similar structure but so far poor experimental data we give the corresponding formulae, to be used in the future.

1 Introduction

The $MSSM+R_p$ is obtained from the $MSSM$ by adding the following terms to the superpotential (*c.f.* ref.[1])

$$\begin{aligned} \Delta\mathcal{W}_{R_p} = & \frac{1}{2} \varepsilon^{ab} \lambda_{ijk} L_a^i L_b^j E^{kC} + \varepsilon^{ab} \delta^{xy} \lambda'_{ijk} L_a^i Q_{bx}^j D_y^{kC} \\ & + \frac{1}{2} \varepsilon^{xyz} \lambda''_{ijk} U_x^{iC} D_y^{jC} D_z^{kC} + \varepsilon^{ab} \kappa_i L_a^i H_b^U. \end{aligned} \quad (1)$$

H, Q, L represent the leftchiral $SU(2)_W$ -doublet superfields of the Higgses, the quarks and leptons; U, D, E represent the rightchiral superfields of the u -type quarks, d -type

*dreiner@th.physik.uni-bonn.de

†polesell@hawk.pv.infn.it

‡thor@thphys.ox.ac.uk

quarks and electron-type leptons, respectively; a superscript C denotes charge conjugation; a, b and x, y, z are $SU(2)_W$ - and $SU(3)_C$ -indices, respectively; i, j, k and later also f, g, l, n are generational indices (summation over repeated indices is implied); δ^{xy} is the Kronecker symbol, $\varepsilon^{\dots}$ symbolizes any tensor that is totally antisymmetric with respect to the exchange of any two indices, with $\varepsilon^{12\dots} = 1$. The coupling constants $\lambda_{ijk}/\lambda''_{ijk}$ are antisymmetric with respect to the exchange of the first two/last two indices. The last term in eq.(1) can be rotated away utilizing a unitary field-redefinition.

Good agreement between SM theory and experiment gives stringent upper bounds on the extra 45 coupling constants λ_{ijk} , λ'_{ijk} and λ''_{ijk} , as well as on products thereof. For a list of references and the processes dealt with, see e.g. ref.[2, 3, 4]. In particular, in $\mathcal{R}_p$ there are new operators for *leptonic* meson decays. The SM theoretical predictions for the decay widths of mesons and the measured values match up within the experimental uncertainty. We can thus determine yet further tight constraints on several products of coupling constants: $\lambda^*\lambda'$ and $\lambda^*\lambda$. This was first done in ref.[5] for single coupling constants and later in ref.[6] for some products, however only treating charged pions decaying via either d -type squark or slepton exchange, respectively. Ref.[7] treated general leptoquark reactions of several particles, one of them the K_L^0 ; this result was quoted in terms of $\mathcal{R}_p$ by ref.[8]; the same result was reached by ref.[9]. Ref.[10] among other things dealt with the decay of K_L^0 , however with only u -squark exchange contributing to SM -allowed processes. The decays of neutral and charged B -mesons were treated in ref.[11] and ref.[12], respectively. We generalize these calculations, focusing on products of two coupling constants, and stress where we obtain new or stricter bounds.

2 $\mathcal{R}_p$ -Decay of Charged Mesons

2.1 Calculation of the Decay Rate

Consider a negatively charged meson π^{ij} at rest made of a d -type quark d^i and a u -type antiquark u^{jC} which decays into an antineutrino ν^{nC} and a charged lepton ℓ^f , i.e.

$$\left| \pi^{ij}(p_1) \right\rangle \longrightarrow \left| \nu^{nC}(p_2) ; \ell^f(p_3) \right\rangle, \quad (2)$$

the $p_{1,2,3}$ being four-momenta. We now calculate the partial decay rate of this process. Focusing on the Yukawa-couplings of the first two terms in eq.(1) leads to, again with summation over repeated indices implied,

$$\mathcal{L}_{\mathcal{R}_p} \supset \lambda_{ijk} \left(\overline{\nu^{iC}} P_L \ell^j \widetilde{\ell_R^{k*}} + \overline{\ell^k} P_L \ell^j \widetilde{\nu_L^i} + \overline{\ell^k} P_L \nu^i \widetilde{\ell_L^j} \right)$$




Figure 1: The tree-level $MSSM+R_p$ processes contributing to the decay of the charged mesons.

$$\begin{aligned}
& + \lambda'_{ijk} \left(\overline{\nu^{iC}} P_L d^j \widetilde{d_R^k}^* + \overline{d^k} P_L d^j \widetilde{\nu_L^i} + \overline{d^k} P_L \nu^i \widetilde{d_L^j} \right) \\
& - \lambda'_{ijk} \left(\overline{u^{jC}} P_L \ell^i \widetilde{d_R^k}^* + \overline{d^k} P_L u^j \widetilde{\ell_L^i} + \overline{d^k} P_L \ell^i \widetilde{u_L^j} \right) + c.c. \quad (3)
\end{aligned}$$

All spinors are Dirac spinors, the overbar denotes the Dirac adjoint, $P_{L,R}$ are the projection operators on the left-/right-handed parts. *The fermions are mass-eigenstates.* A tilde denotes a scalar; the scalars' subscripts L, R indicate the chirality of the corresponding Weyl spinor. The 4th term in eq.(3) together with the *c.c.* of the 7th term, and the 3rd term together with the *c.c.* of the 8th term lead to the meson decay processes depicted in fig. 1, which give the effective Hamiltonians

$$\begin{aligned}
\mathcal{H}^{\widetilde{d_R}} &= \frac{1}{2} \sum_k \frac{\lambda_{fjk}^* \lambda'_{nik}}{m_{\widetilde{d_R^k}}^2} \overline{\ell^f} \gamma_\nu P_L \nu^n \overline{u^j} \gamma^\nu P_L d^i, \\
\mathcal{H}^{\widetilde{\ell_L}} &= - \sum_k \frac{\lambda_{kji}^* \lambda_{nkf}}{m_{\widetilde{\ell_L^k}}^2} \overline{\ell^f} P_L \nu^n \overline{u^j} P_R d^i, \quad (4)
\end{aligned}$$

where m is the mass of a particle. To obtain the first equation we employed a Fierz-identity. These two Hamiltonians have to be added to the effective Hamiltonian for the SM -process

$$\mathcal{H}^W = \frac{4 G_F V_{ji}}{\sqrt{2}} \overline{\ell^f} \gamma_\nu P_L \nu^f \overline{u^j} \gamma^\nu P_L d^i. \quad (5)$$

Here G_F is the Fermi constant and V_{ji} is an element of the CKM-matrix. We obtain for the transition amplitude $\mathcal{M}_{ijfn}$

$$\mathcal{M}_{ijfn} \delta^4(p_1 - p_2 - p_3) = \frac{1}{2\pi i} \int \langle \ell^f ; \nu^{nC} | (\mathcal{H}^W + \mathcal{H}^{\widetilde{d_R}} + \mathcal{H}^{\widetilde{\ell_L}}) | \pi^{ij} \rangle d^4x. \quad (6)$$

We expand the fields in the initial and final state, perform the integrations and use

$$\begin{aligned} \langle 0 | \overline{u^j}(y) \gamma^\nu P_{L,R} d^i(y) | \pi^{ij}(p_1) \rangle &= \pm \frac{1}{\sqrt{2}} f_{\pi^{ij}} p_1^\nu e^{-ip_1 y}, \\ \langle 0 | \overline{u^j}(y) P_{L,R} d^i(y) | \pi^{ij}(p_1) \rangle &= \mp \frac{1}{\sqrt{2}} \frac{m_{\pi^{ij}}^2}{m_{u^j} + m_{d^i}} f_{\pi^{ij}} e^{-ip_1 y}; \end{aligned} \quad (7)$$

$f_{\pi^{ij}}$ is the meson decay constant.¹ Thus

$$\begin{aligned} \mathcal{M}_{ijfn} &= \frac{f_{\pi^{ij}}}{2\sqrt{2}} \overline{\mathcal{U}}^f(\vec{p}_3) \times \\ &\sum_k \left\{ \left(\frac{\delta_{fn}}{3} \frac{8 G_F V_{ji}}{\sqrt{2}} + \frac{\lambda_{fjk}^* \lambda'_{nik}}{m_{d_R^k}^2} \right) \not{p}_1 - 2 \frac{m_{\pi^{ij}}^2}{m_{u^j} + m_{d^i}} \frac{\lambda_{kji}^* \lambda_{nkf}}{m_{\ell_L^k}^2} \right\} \\ &\times P_L \mathcal{V}^n(\vec{p}_2). \end{aligned} \quad (8)$$

$\overline{\mathcal{U}}^f, \mathcal{V}^n$ are the Fourier coefficient functions of ℓ^f, ν^{nC} , respectively. Next we take the absolute value square, average over the spins and use the trace theorems. Then we sum over n , because the experiments that measured the partial decay widths did not determine the flavour of the antineutrinos,² resulting in

$$\sum_n \langle |\mathcal{M}_{ijfn}|^2 \rangle = 4G_F^2 f_{\pi^{ij}}^2 |V_{ji}|^2 (m_{\pi^{ij}}^2 - m_{\ell^f}^2) m_{\ell^f}^2 \sum_n \left(\delta_{fn} + 2\delta_{fn} \text{Re}[K_{ijfn}] + |K_{ijfn}|^2 \right), \quad (9)$$

with

$$K_{ijfn} = \frac{\sqrt{2}}{8 G_F |V_{ji}|} \sum_k \left(\frac{\lambda_{fjk}^* \lambda'_{nik}}{m_{d_R^k}^2} - 2 \frac{m_{\pi^{ij}}^2}{m_{\ell^f} (m_{u^j} + m_{d^i})} \frac{\lambda_{kji}^* \lambda_{nkf}}{m_{\ell_L^k}^2} \right), \quad (10)$$

containing all $\mathcal{R}_p$ contributions; $2\text{Re}[K_{ijfn}]$ in eq.(9) is due to the interference between SM and $\mathcal{R}_p$ amplitudes. For simplicity we neglect the phase of the CKM-matrix. The partial decay rate is then

$$\Gamma_{\pi^{ij} \rightarrow \ell^f \nu^{fC}}^{SM+\mathcal{R}_p} = \Gamma_{\pi^{ij} \rightarrow \ell^f \nu^{fC}}^{SM} \times \left(1 + 2\text{Re}[K_{ijff}] + \sum_n |K_{ijfn}|^2 \right), \quad (11)$$

with ν^C being an *arbitrary* antineutrino, and

$$\Gamma_{\pi^{ij} \rightarrow \ell^f \nu^{fC}}^{SM} = C_{ijf} G_F^2 f_{\pi^{ij}}^2 |V_{ji}|^2 \frac{((m_{\pi^{ij}}^2 - m_{\ell^f}^2) m_{\ell^f})^2}{4\pi m_{\pi^{ij}}^3}; \quad (12)$$

the correction factor C_{ijf} of $\mathcal{O}(1)$ is due to higher order electroweak leading logarithms, short distance QCD corrections, and structure dependent effects, see ref.[13] and also ref.[14].

¹There are several ways of defining the meson decay constant, differing by factors of $\sqrt{2}$; in the convention we use $f_\pi = (92.4 \pm 0.3) \text{ MeV}$, see ref.[15].

²The upper experimental bounds on $\pi^- \rightarrow \mu \nu_e^C$ and $K^- \rightarrow \mu \nu_e^C$, see ref.[15], come from a different type of experiment, compared to the one used to determine the branching ratios for $\pi^- \rightarrow \mu \nu^C$ and $K^- \rightarrow \mu \nu^C$. They do not lead to better bounds on the coupling constants.

2.2 Calculation of the Bounds

We prefer not to compare the experimental data directly with eq.(11), since $f_{\pi^{ij}}$ has quite a large error. This leads to very weak bounds on K_{ijfn} . To avoid this, we introduce

$$\mathcal{R}_{\pi^{ij}} := \frac{\Gamma_{\pi^{ij} \rightarrow \ell f \nu^C}}{\Gamma_{\pi^{ij} \rightarrow \ell^g \nu^C}}, \quad (13)$$

with $m_{\ell^g} > m_{\ell f}$. If the experimental and SM -theoretical decay rates agree well we have, see eq.(11), $|2\text{Re}[K_{ijff}] + \sum_n |K_{ijfn}|^2| \ll 1$. Plugging eq.(11) into eq.(13) one gets

$$\frac{\mathcal{R}_{\pi^{ij}}^{SM+\mathcal{R}_p}}{\mathcal{R}_{\pi^{ij}}^{SM}} := 1 + \epsilon_{\pi^{ij}} \approx 1 + 2 \text{Re}[K_{ijff} - K_{ijgg}] + \sum_n |K_{ijfn}|^2 - \sum_n |K_{ijgn}|^2. \quad (14)$$

Let $\Delta \dots$ symbolize the theoretical or experimental uncertainty. If the theoretical prediction $\mathcal{R}_{\pi^{ij}}^{SM} \pm \Delta \mathcal{R}_{\pi^{ij}}^{SM}$ lies within the experimental range $\mathcal{R}_{\pi^{ij}}^{exp} \pm \Delta \mathcal{R}_{\pi^{ij}}^{exp}$, one has

$$\epsilon_{\pi^{ij}}^{min} := \frac{\mathcal{R}_{\pi^{ij}}^{exp}}{\mathcal{R}_{\pi^{ij}}^{SM}} - \Delta \left(\frac{\mathcal{R}_{\pi^{ij}}^{exp}}{\mathcal{R}_{\pi^{ij}}^{SM}} \right) - 1 \leq \epsilon_{\pi^{ij}} \leq \frac{\mathcal{R}_{\pi^{ij}}^{exp}}{\mathcal{R}_{\pi^{ij}}^{SM}} + \Delta \left(\frac{\mathcal{R}_{\pi^{ij}}^{exp}}{\mathcal{R}_{\pi^{ij}}^{SM}} \right) - 1 =: \epsilon_{\pi^{ij}}^{max}. \quad (15)$$

We could use this to determine a bound on this general combination of $\mathcal{R}_p$ coupling constants; however, the bounds on individual coupling constants are typically of the order $\mathcal{O}(10^{-2})$, see ref.[2], and thus we limit ourselves to at most two non-zero coupling constants at a time, and in each case suppose the other 34 λ, λ' coupling constants vanish (eq.(16), eq.(17) and eq.(18) are also valid for $f \rightarrow g$ with $\epsilon_{\pi^{ij}}^{max} \leftrightarrow -\epsilon_{\pi^{ij}}^{min}$):

$$\epsilon_{\pi^{ij}}^{min} \leq 2 \text{Re}[K_{ijff}] + |K_{ijff}|^2 \leq \epsilon_{\pi^{ij}}^{max} \quad \text{and for } n \neq f \quad |K_{ijfn}|^2 \leq \epsilon_{\pi^{ij}}^{max}. \quad (16)$$

We assume that the imaginary parts of the coupling constants are approximately the same as the corresponding real parts.³ With $G_F = (0.116639 \pm 0.000001) \times (100 \text{ GeV})^{-2}$, see ref.[15], we obtain

$$\begin{aligned} -0.330 |V_{ji}| \left(\sqrt{1 + 2 \epsilon_{\pi^{ij}}^{min}} + 1 \right) &\leq \\ &\frac{\text{Re}[\lambda'_{fjk}^* \lambda'_{fik}]}{(m_{\widetilde{d_R^k}}/100 \text{ GeV})^2}, \quad \frac{-2 m_{\pi^{ij}}^2}{m_{\ell f}(m_{u^j} + m_{d^i})} \frac{\text{Re}[\lambda'_{kji}^* \lambda_{fkf}]}{(m_{\widetilde{\ell_L^k}}/100 \text{ GeV})^2} \\ &\leq 0.330 |V_{ji}| \left(\sqrt{1 + 2 \epsilon_{\pi^{ij}}^{max}} - 1 \right) \end{aligned} \quad (17)$$

and for $n \neq f$

$$\frac{|\lambda'_{fjk}^* \lambda'_{nik}|}{(m_{\widetilde{d_R^k}}/100 \text{ GeV})^2}, \quad \frac{2 m_{\pi^{ij}}^2}{m_{\ell f}(m_{u^j} + m_{d^i})} \frac{|\lambda'_{kji}^* \lambda_{nkf}|}{(m_{\widetilde{\ell_L^k}}/100 \text{ GeV})^2} \leq 0.66 |V_{ji}| \sqrt{\epsilon_{\pi^{ij}}^{max}}. \quad (18)$$

The prefactor $2m_{\pi^{ij}}^2/[m_{\ell f}(m_{u^j} + m_{d^i})]$ results in much tighter bounds for $\lambda'_{kji}^* \lambda_{nkf}$. We will apply these results only to processes with sufficiently small experimental error bars.

³If the imaginary part vanishes the bounds are weaker by a factor of $\mathcal{O}(1)$.

2.3 $\pi^- \rightarrow \ell^{f,g} + \nu^C$

As a first application, we consider pion decay with $f, i, j = 1, g = 2$. The SM gives the 2σ theoretical value $\mathcal{R}_{\pi^-}^{SM} = (1.2354 \pm 0.0004) \times 10^{-4}$ (see ref.[14]; the uncertainty mainly derives from C_{111} and C_{112}). From the partial decay widths at the 2σ level in ref.[15], namely $\Gamma_{\pi^- \rightarrow e\nu^C}^{exp}/\Gamma_{\pi^-}^{exp\ total} = (1.230 \pm 0.008) \times 10^{-4}$ and $\Gamma_{\pi^- \rightarrow \mu\nu^C}^{exp}/\Gamma_{\pi^-}^{exp\ total} = 0.9998770 \pm 8 \times 10^{-7}$, one calculates $\mathcal{R}_{\pi^-}^{exp} = (1.230 \pm 0.008) \times 10^{-4}$. Hence, $\epsilon_{\pi^-}^{min} = -0.0107$ and $\epsilon_{\pi^-}^{max} = 0.0022$. With $|V_{11}| = 0.9750 \pm 0.0008$, ref.[5] obtained bounds on a single coupling constant; this was updated in ref.[16]. We have reproduced their results. The experimental data have only marginally changed and the new bounds are $|\lambda'_{11k}| \leq 0.027 \ m_{d_R^k}/100 \text{ GeV}$ and $|\lambda'_{21k}| \leq 0.059 \ m_{d_R^k}/100 \text{ GeV}$. We obtain bounds for the products of couplings $|\lambda'_{11k} \lambda'_{21k}| \leq 0.03 (m_{d_R^k}/100 \text{ GeV})^2$, $|\lambda'_{11k} \lambda'_{31k}| \leq 0.03 (m_{d_R^k}/100 \text{ GeV})^2$ and $|\lambda'_{21k} \lambda'_{31k}| \leq 0.066 (m_{d_R^k}/100 \text{ GeV})^2$. The first bound is redundant since the product of the single bounds is stronger; the second and the third bound are almost the same as the single bound on $|\lambda'_{11k}|$ and $|\lambda'_{21k}|$. Furthermore, we obtain the following new bounds using $m_e = (0.510998902 \pm 2.1 \times 10^{-8}) \text{ MeV}$, $m_\mu = (105.6583568 \pm 5.2 \times 10^{-6}) \text{ MeV}$, $m_{\pi^-} = (139.57018 \pm 0.00035) \text{ MeV}$ and $m_u + m_d = (8.5 \pm 3.5) \text{ MeV}$,⁴ see ref.[15],

$$\begin{aligned}
-7.9 \times 10^{-8} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2 &\leq \text{Re}[\lambda'_{k11} \lambda_{1k1}] \leq 7.1 \times 10^{-5} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2, \\
-7.9 \times 10^{-5} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2 &\leq \text{Re}[\lambda'_{311} \lambda_{232}] \quad , \\
|\lambda'_{k11} \lambda_{3k1}| &\leq 3.4 \times 10^{-6} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda'_{211} \lambda_{322}| &\leq 1.5 \times 10^{-3} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda'_{111} \lambda_{211}| &\leq 3.4 \times 10^{-6} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2. \quad (19)
\end{aligned}$$

The upper bound/bounds we obtained for $\text{Re}[\lambda'_{311} \lambda_{232}]/\text{Re}[\lambda'_{111} \lambda_{212}]$, $|\lambda'_{111} \lambda_{312}|$ are weaker than the products of the two bounds on the single coupling constants, see ref.[4]; there also much stricter bounds were stated for $|\lambda'_{k11} \lambda_{1k2}|$ as well as for $|\lambda'_{311} \lambda_{231}|$.

2.4 $K^- \rightarrow \ell^{f,g} + \nu^C$

Next we consider charged kaon decay with $f, j = 1, g, i = 2$. According to ref.[14], $\mathcal{R}_{K^-}^{SM} = (2.472 \pm 0.002) \times 10^{-5}$ at the 2σ level. Experimentally $\Gamma_{K^- \rightarrow e\nu^C}^{exp}/\Gamma_{K^-}^{exp\ total} =$

⁴This and m_s are the biggest sources of inaccuracy, going linearly into the bounds on $\lambda'^* \lambda$.

$(1.55 \pm 0.14) \times 10^{-5}$ and $\Gamma_{K^- \rightarrow \mu \nu^C}^{exp} / \Gamma_{K^-}^{exp total} = 0.6351 \pm 0.0036$, at the 2σ level [15]. Therefore $\mathcal{R}_{K^-}^{exp} = (2.44 \pm 0.22) \times 10^{-5}$ and $\epsilon_{K^-}^{min} = -0.10$ and $\epsilon_{K^-}^{max} = 0.076$. Using $|V_{12}| = 0.222 \pm 0.004$ we obtain $|\lambda_{11k}^* \lambda'_{32k}| \leq 0.04 (m_{\widetilde{d_R^k}}/100 \text{ GeV})^2$, $|\lambda_{21k}^* \lambda'_{32k}| \leq 0.046 (m_{\widetilde{d_R^k}}/100 \text{ GeV})^2$. As for the pion these bounds are almost the same as the ones on $|\lambda'_{11k}|$, $|\lambda'_{21k}|$. Our bounds on $\text{Re}[\lambda_{11k}^* \lambda'_{12k}]$, $\text{Re}[\lambda_{21k}^* \lambda'_{22k}]$ are much weaker than the bounds on $|\lambda_{i1k}^* \lambda'_{i2k}|$, see ref.[4], and we do not list them. Similarly, the existing bounds on $|\lambda_{11k}^* \lambda'_{22k}|$, $|\lambda_{21k}^* \lambda'_{12k}|$ are much stronger than ours. Furthermore with $m_{K^-} = (493.677 \pm 0.016) \text{ MeV}$, $m_s = (122.5 \pm 47.5) \text{ MeV}$ and $m_s = (21 \pm 4)m_d$ (see ref.[15]) we have the following new bounds

$$\begin{aligned}
-7.0 \times 10^{-7} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2 &\leq \text{Re}[\lambda_{k12}^* \lambda_{1k1}] \leq 1.8 \times 10^{-5} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2, \\
-1.8 \times 10^{-4} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2 &\leq \text{Re}[\lambda_{k12}^* \lambda_{2k2}] \leq 3.8 \times 10^{-3} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2 \quad (k=3), \\
|\lambda_{k12}^* \lambda_{2k1}| &\leq 5.4 \times 10^{-6} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k12}^* \lambda_{3k1}| &\leq 5.4 \times 10^{-6} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k12}^* \lambda_{1k2}| &\leq 1.3 \times 10^{-3} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k12}^* \lambda_{3k2}| &\leq 1.3 \times 10^{-3} \left(\frac{m_{\widetilde{\ell_L^k}}}{100 \text{ GeV}} \right)^2. \tag{20}
\end{aligned}$$

The upper bound on $\text{Re}[\lambda_{k12}^* \lambda_{212}]$ obtained from two bounds on the single coupling constants is stricter than the one we obtained.

2.5 $B^- \rightarrow \ell^f + \nu^C$

For the charged B -meson decay the procedure is slightly different since it has not been directly measured. Unlike the two previous cases one only has an experimental upper bound on the branching ratio $\mathcal{B}$, see ref.[15], and thus has to go back to eq.(11). This has been done in ref.[12]. We go beyond their work with a more conservative account of the experimental errors and obtain weaker bounds. We also work from the beginning in the mass eigenstate basis to avoid model dependent results, see ref.[18].

First $f=3$. The theoretical predictions are limited by $\Gamma_{B \rightarrow \tau \nu^C}^{SM+\mathcal{R}_p} / \Gamma_B^{SM+\mathcal{R}_p total} \leq 5.7 \times 10^{-4}$. As the total widths $\Gamma_B^{exp total}$ and $\Gamma_B^{SM total}$ agree fairly well one has $\Gamma_B^{SM+\mathcal{R}_p total} \approx \Gamma_B^{SM total}$, so

that, utilizing eq.(11), we obtain for the branching ratio

$$\frac{\Gamma_{B \rightarrow \tau \nu^C}^{SM+\mathcal{R}_p}}{\Gamma_{B \text{ total}}^{SM+\mathcal{R}_p}} \approx \left(1 + 2 \operatorname{Re}[K_{3133}] + \sum_n |K_{313n}|^2\right) \frac{\Gamma_{B \rightarrow \tau \nu^C}^{SM}}{\Gamma_{B \text{ total}}^{SM}}. \quad (21)$$

To keep the combined uncertainties of $|V_{13}|$ and f_B as small as possible we use the theoretical prediction (see ref.[17])

$$\frac{\Gamma_{B \rightarrow \tau \nu^C}^{SM}}{\Gamma_{B \text{ total}}^{SM}} = (4.08 \pm 0.24) \times 10^{-4} \left| \frac{V_{13}}{V_{31}} \right|^2. \quad (22)$$

In order to take into account the correlated uncertainties in V_{13}/V_{31} we use the Wolfenstein parameterization (see e.g. ref.[19]):

$$\frac{V_{13}}{V_{31}} = \frac{\bar{\rho} - i\bar{\eta}}{1 - \frac{\lambda^2}{2} - \bar{\rho} - i\bar{\eta}}. \quad (23)$$

The Wolfenstein parameters are given by (see ref.[20]) $\bar{\rho} = 0.21 \pm 0.12$, $\bar{\eta} = 0.38 \pm 0.11$, $\lambda = 0.222 \pm 0.004$, all at 95% C.L.. We thus obtain for the theoretical prediction $\Gamma_{B \rightarrow \tau \nu^C}^{SM}/\Gamma_{B \text{ total}}^{SM} = (1.05 \pm 0.65) \times 10^{-4}$. The lower value should be used in eq.(21), to be compared with the experimental upper bound. Thus

$$2 \operatorname{Re}[K_{3133}] + \sum_n |K_{313n}|^2 \leq 13.3. \quad (24)$$

In the following, we again assume that only two coupling constants are non-zero. Thus we have $|K_{3131}|, |K_{3132}| \leq \sqrt{13.3}$. Furthermore the imaginary part is again taken to be about the same as the real part, hence $-\sqrt{\frac{1}{4} + \frac{13.3}{2}} - \frac{1}{2} \leq \operatorname{Re}[K_{3133}] \leq \sqrt{\frac{1}{4} + \frac{13.3}{2}} - \frac{1}{2}$. Thus, with $|V_{13}| = 0.0035 \pm 0.0015$ (see ref.[15]), $m_{B^-} = (5279.0 \pm 0.5) \text{ MeV}$, $m_b = (4200 \pm 200) \text{ MeV}$ and $m_\tau = (1776.99 \pm 0.29) \text{ MeV}$, see ref.[15], we obtain

$$\begin{aligned} |\lambda_{313}^* \lambda_{233}| &\leq 2 \times 10^{-3} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2, \\ -6 \times 10^{-4} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2 &\leq \operatorname{Re}[\lambda_{213}^* \lambda_{323}] \leq 1 \times 10^{-3} \left(\frac{m_{\ell_L^k}}{100 \text{ GeV}} \right)^2. \end{aligned} \quad (25)$$

According to ref.[12] the bounds on $|\lambda_{k13}^* \lambda_{1k3}|$, $|\lambda_{31k}^* \lambda'_{13k}|$, $|\lambda_{31k}^* \lambda'_{23k}|$, $\operatorname{Re}[\lambda_{31k}^* \lambda'_{33k}]$ and $\operatorname{Re}[\lambda_{113}^* \lambda_{313}]$ are not better than the previous ones; furthermore the bound on $|\lambda_{113}^* \lambda_{213}|$ is weaker than the product of the two bounds on the single coupling constants, see ref.[4].

Analogously, for $f = 1, 2$,

$$\left(1 + 2 \operatorname{Re}[K_{31ff}] + \sum_n |K_{31fn}|^2\right) \Gamma_{B \rightarrow \ell^f \nu^f C}^{SM} \times \tau_{B^-} < 5.7 \times 10^{-4}, \quad (26)$$

where τ_{B^-} is the B -meson life time. Instead of arguing that the error on $\Gamma_{B \rightarrow \ell^f \nu^f C}^{SM}$ is $\pm \frac{0.65}{1.05} \Gamma_{B \rightarrow \ell^f \nu^f C}^{SM}$, see the third line below eq.(23), we are going to be as conservative as

possible. Due to isospin invariance B^0 and B^- have the same decay constant. From ref.[21], $f_{B^0} = (200 \pm 30)$ MeV, and thus with our convention $f_B = (141 \pm 21)$ MeV (*c.f.* footnote 1). Therefore, $f_B^2 |V_{13}|^2 = (0.24 \pm 0.22) \text{ MeV}^2$, and with $\tau_{B^-} = 1.655 \times 10^{-12}$ s (see ref.[15]), we obtain for $f = 1$ that $(1 + \dots)(9.0 \pm 8.3) \times 10^{-12} \leq 1.5 \times 10^{-5}$ and for $f = 2$ that $(1 + \dots)(3.8 \pm 3.5) \times 10^{-7} \leq 2.1 \times 10^{-5}$. Working with the lower value, for $f = 1$ we get

$$|\lambda'_{k13} \lambda_{3k1}| \leq 6 \times 10^{-4} \left(\frac{m_{\widetilde{\ell}_L^k}}{100 \text{ GeV}} \right)^2. \quad (27)$$

The bounds on $|\lambda'_{k13} \lambda_{2k1}|$ and $\text{Re}[\lambda'_{k13} \lambda_{1k1}]$ are not improved compared to the previous ones, see ref.[12], and the ones on $|\lambda'_{11k} \lambda'_{23k}|$, $|\lambda'_{11k} \lambda'_{33k}|$ and $\text{Re}[\lambda'_{11k} \lambda'_{13k}]$ are too poor to be listed. $f = 2$ yields

$$|\lambda'_{k13} \lambda_{3k2}| \leq 7 \times 10^{-4} \left(\frac{m_{\widetilde{\ell}_L^k}}{100 \text{ GeV}} \right)^2. \quad (28)$$

Ref.[12] states that there exist better bounds on $|\lambda'_{k13} \lambda_{1k2}|$ and $\text{Re}[\lambda'_{k13} \lambda_{2k2}]$. The bounds on $|\lambda'_{21k} \lambda'_{13k}|$, $|\lambda'_{21k} \lambda'_{33k}|$, $\text{Re}[\lambda'_{21k} \lambda'_{23k}]$ are almost the same as the single bound on $|\lambda'_{21k}|$.

3 $\widetilde{R}_p$ -Decay of Neutral Mesons

3.1 Calculation of the Bounds

Now we deal with the bound state $(d^j C d^i)$ decaying into ℓ^f and ℓ^{nC} , with momenta p_1, p_3, p_2 , respectively. We consider only $n \neq f$, in which case the process does not occur in the SM and therefore no contributions from loop-diagrams have to be taken into account. We proceed as in the previous section. In eq.(3) the 9^{th} term together with its *c.c.* contributes to the decay, in analogy to the $\widetilde{d_R^k}$ -exchange in the last section. Furthermore the 2^{nd} term together with the *c.c.* of the 5^{th} and the 5^{th} term together with the *c.c.* of the 2^{nd} contribute, both in analogy to the $\widetilde{\ell_L^k}$ -exchange. The Hamiltonian is given by

$$\begin{aligned} \mathcal{H}_{\widetilde{u_L}} &= \frac{1}{2} \sum_k \frac{\lambda'_{nkj} \lambda_{fki}^*}{m_{\widetilde{u_L^k}}^2} \overline{\ell^f} \gamma^\nu P_L \ell^n \overline{d^j} \gamma_\nu P_R d^i, \\ \mathcal{H}_{\widetilde{\nu_L}} &= \sum_k \frac{\lambda_{kji}^* \lambda_{knf}}{m_{\widetilde{\nu_L^k}}^2} \overline{\ell^f} P_L \ell^n \overline{d^j} P_R d^i \\ &+ \sum_k \frac{\lambda'_{kij} \lambda_{kfn}^*}{m_{\widetilde{\nu_L^k}}^2} \overline{\ell^f} P_R \ell^n \overline{d^j} P_L d^i. \end{aligned} \quad (29)$$

Using the results corresponding to eq.(7) we obtain

$$\mathcal{M}_{ijfn} = -\frac{f_{(d^j C d^i)}}{2\sqrt{2}} \overline{\mathcal{U}^f}(\vec{p}_3) \left\{ A_{ijfn} \not{p}_1 P_L + B_{ijfn} P_L - B_{jinf}^* P_R \right\} \mathcal{V}^n(\vec{p}_2), \quad (30)$$

where

$$A_{ijfn} = \sum_k \frac{\lambda'_{nkj} \lambda_{fki}^*}{m_{\widetilde{u_L^k}}^2} = A_{jinf}^*, \quad B_{ijfn} = 2 \frac{m_{(d^j C d^i)}^2}{(m_{d^j} + m_{d^i})} \sum_k \frac{\lambda_{kji}^* \lambda_{knf}}{m_{\widetilde{\nu_L^k}}^2}. \quad (31)$$

Hence

$$\begin{aligned} \Gamma_{(d^j C d^i) \rightarrow \ell^f + \ell^n C}^{SM+\mathcal{R}_p} &= \sqrt{m_{(d^j C d^i)}^4 + m_{\ell^f}^4 + m_{\ell^n}^4 - 2(m_{(d^j C d^i)}^2 m_{\ell^f}^2 + m_{\ell^f}^2 m_{\ell^n}^2 + m_{\ell^n}^2 m_{(d^j C d^i)}^2)} \\ &\times \frac{f_{(d^j C d^i)}^2}{128 \pi m_{(d^j C d^i)}^3} \times \Upsilon_{ijfn}, \end{aligned} \quad (32)$$

where

$$\begin{aligned} \Upsilon_{ijfn} &= (m_{(d^j C d^i)}^2 - m_{\ell^f}^2) |A_{ijfn} m_{\ell^f} + B_{ijfn}|^2 + (m_{(d^j C d^i)}^2 - m_{\ell^n}^2) |A_{ijfn} m_{\ell^n} + B_{jinf}^*|^2 \\ &- |B_{ijfn} m_{\ell^n} - B_{jinf}^* m_{\ell^f}|^2 + m_{\ell^f} m_{\ell^n} \left\{ |B_{ijfn} + B_{jinf}^*|^2 - |A_{ijfn} m_{\ell^n} - B_{ijfn}|^2 \right. \\ &\left. - |A_{ijfn} m_{\ell^f} - B_{jinf}^*|^2 + |(m_{\ell^f} + m_{\ell^n}) A_{ijfn}|^2 \right\}. \end{aligned} \quad (33)$$

Due to the large experimental error in $f_{(d^j C d^i)}$, we can neglect m_{ℓ^n} compared to m_{ℓ^f} (with f, n chosen correspondingly) and m_{ℓ^f} compared to $m_{(d^j C d^i)}$. Thus, focusing again on the bounds on products of two coupling constants, with all other coupling constants vanishing,

$$|A_{ijfn}|, \frac{|B_{ijfn}|}{m_{\ell^f}}, \frac{|B_{jinf}^*|}{m_{\ell^f}} \leq \frac{20}{f_{(d^j C d^i)} m_{\ell^f}} \sqrt{\frac{\Gamma_{(d^j C d^i) \rightarrow \ell^f + \ell^n C}^{exp. upper bound} / \Gamma_{(d^j C d^i) \rightarrow total}^{exp}}{m_{(d^j C d^i)} \tau_{(d^j C d^i)}^{exp}}}. \quad (34)$$

Here τ is the mean life time. The same considerations apply to mesons that have wave functions of the form

$$\pi_{ij}^0 = \frac{1}{\sqrt{2}} [(d^j C d^i) \pm (d^i C d^j)]; \quad (35)$$

one replaces every A_{ijfn} by $\frac{1}{\sqrt{2}}(A_{ijfn} \pm A_{jinf})$, and likewise for B_{ijfn} , B_{jinf}^* . As in the previous section, we will apply eq.(34) only to processes with satisfactory experimental data, as was done in ref.[11], treating among other processes $B^0 \rightarrow \ell^f + \ell^n C$; we confirm their results.

3.2 $B_s^0 \rightarrow \mu + e^C$

We now consider $B_s^0, B_s^{0C} \rightarrow \mu + e^C$, $i, f = 2, j = 3, n = 1$ and $i \leftrightarrow j$. The relevant parameters are given by $f_{B_s^0} = (1.16 \pm 0.04) f_{B^0}$, see ref.[21], $\mathcal{B}(B_s^0 \rightarrow \mu + e^C) < 6.1 \times 10^{-6}$, $\tau_{B_s^0} = (1.464 \pm 0.057) \times 10^{-12}$ s, $m_{B_s^0} = (5369.6 \pm 2.4)$ MeV, see ref.[15]. Thus

$$\begin{aligned}
|\lambda'_{123} \lambda_{222}^*| &\leq 8 \times 10^{-3} \left(\frac{m_{u_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda'_{1k2} \lambda_{2k3}^*| &\leq 8 \times 10^{-3} \left(\frac{m_{u_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k32}^* \lambda_{k12}| &\leq 7 \times 10^{-5} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k23}^* \lambda_{k21}| &\leq 7 \times 10^{-5} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k32}^* \lambda_{k21}| &\leq 7 \times 10^{-5} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k23}^* \lambda_{k12}| &\leq 7 \times 10^{-5} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2.
\end{aligned} \tag{36}$$

Our results for $|\lambda'_{1k3} \lambda_{2k2}^*|$ ($k \neq 2$) and $k = 1$ for the second bound are weaker than the products of the bounds on single coupling constants, see ref.[4].

3.3 $K_L^0 \rightarrow \mu + e^C$

K_L^0 is defined as $[K_2^0 + \epsilon K_1^0]/\sqrt{1 + \epsilon^2}$, with $K_{1,2}^0 = [K^0 \pm K^{0C}]/\sqrt{2}$. ϵ parameterizes the CP -violation. If we neglect ϵ , $K_L^0 = [K^0 - K^{0C}]/\sqrt{2}$, with $K^0 = (s^C d)$. From ref.[15] one has $m_{K_L^0} = (497.672 \pm 0.031)$ MeV, $\tau_{K_L^0} = (5.17 \pm 0.04) \times 10^{-8}$ s and $\mathcal{B}(K_L^0 \rightarrow \mu + e^C) < 4.7 \times 10^{-12}$. Ref.[15] gives $f_K = (159 \pm 1.4 \pm 0.44)$ MeV, which in the convention we use gives the central value 112.4 MeV. Hence, the first two bounds updating previous ones,

$$\begin{aligned}
|\lambda'_{1k2} \lambda_{2k1}^*| &\leq 3 \times 10^{-7} \left(\frac{m_{u_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda'_{1k1} \lambda_{2k2}^*| &\leq 3 \times 10^{-7} \left(\frac{m_{u_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k21}^* \lambda_{k12}| &\leq 6 \times 10^{-9} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda_{k12}^* \lambda_{k12}| &\leq 6 \times 10^{-9} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2, \\
|\lambda'_{k12} \lambda_{k21}^*| &\leq 6 \times 10^{-9} \left(\frac{m_{\nu_L^k}}{100 \text{ GeV}} \right)^2,
\end{aligned}$$

$$|\lambda'_{k21} \lambda^*_{k21}| \leq 6 \times 10^{-9} \left(\frac{m_{\widetilde{\nu_L^k}}}{100 \text{ GeV}} \right)^2. \quad (37)$$

3.4 $\pi^0 \rightarrow \mu + e^C$

With small modifications the result of section 3.1 can also be carried over to admixtures of $(d^j C d^i)$ with $(u^j C u^i)$, as the latter term does not contribute to any decay because the u -type quarks do not couple together to the $\mathcal{R}_p$ operators. However, we shall limit ourselves to the π^0 : η and η' are more complicated, see ref.[15], and the experimental data do not suffice to extract satisfactory bounds.

The relevant parameters here are $m_{\pi^0} = (134.9766 \pm 0.0006) \text{ MeV}$, $\tau_{\pi^0} = (8.4 \pm 0.6) \times 10^{-17} \text{ s}$ and $\mathcal{B}(\pi^0 \rightarrow \mu + e^C) < 3.8 \times 10^{-10}$, see ref.[15]. Thus

$$\begin{aligned} |\lambda'_{311} \lambda^*_{312}| &\leq 3 \times 10^{-3} \left(\frac{m_{\widetilde{\nu_L^k}}}{100 \text{ GeV}} \right)^2, \\ |\lambda'_{311} \lambda^*_{321}| &\leq 3 \times 10^{-3} \left(\frac{m_{\widetilde{\nu_L^k}}}{100 \text{ GeV}} \right)^2. \end{aligned} \quad (38)$$

In ref.[4] a much stronger bound is stated for $|\lambda'_{1k1} \lambda^*_{2k1}|$. Furthermore they present a better bound for $|\lambda'_{111} \lambda^*_{121}|$; and from ref.[4] one finds a stricter bound on $|\lambda'_{211} \lambda^*_{212}|$, based on the bounds on single coupling constants.

4 Summary

We have determined the bounds on products of $\mathcal{R}_p$ coupling constants from leptonic meson decays. In many cases these bounds are better than previous bounds. We have summarized the bounds in the tables at the end of this text. With the formulae given the bounds can easily be updated when the data improve. Furthermore, if additional decays are measured (e.g. from the B -factories) one can determine additional bounds. Eq.(17) and eq.(18) can be used to consider 12 cases: D^- ($i = 1, j = 2$), D_s^- ($i = 2, j = 2$), B^- ($i = 3, j = 1$), B_c^- ($i = 3, j = 2$) decaying into $e + \nu^C$ and $\mu + \nu^C$ ($f = 1, g = 2$), $e + \nu^C$ and $\tau + \nu^C$ ($f = 1, g = 3$), $\mu + \nu^C$ and $\tau + \nu^C$ ($f = 2, g = 3$); eq.(34) can be applied to the decay of B_s^0 ($i = 2, j = 3$) to $\tau + e^C$ ($f = 3, n = 1$) or $\tau + \mu^C$ ($f = 3, n = 2$), and the decay of the Υ , ($i = j = 3$) to $\tau + e^C$ or $\tau + \mu^C$ or $\mu + e^C$ ($f = 2, n = 1$).

5 Acknowledgements

Special thanks to Peter Richardson and Michael Kobel for useful discussions. M.T. would like to thank the Physikalisches Institut, Universität Bonn for their kind hospitality

during most of this work, and the Evangelisches Studienwerk and Worcester College, University of Oxford, for financial support.

References

- [1] S.Weinberg, *Phys.Rev.* **D26** 287 (1982).
- [2] H.Dreiner, *hep-ph/9707435*.
- [3] G.Bhattacharyya, *Nucl.Phys.Proc.Suppl.* **52A** 83 (1997); *hep-ph/9608415*.
- [4] B.Allanach, A.Dedes, H.Dreiner, *Phys.Rev.* **D60** 075014 (1999); *hep-ph/9906209*.
- [5] V.Barger, G.Giudice, T.Han, *Phys.Rev.* **D40** 2987 (1989).
- [6] D.Choudhury, H.Dreiner, P.Richardson, S.Sarkar, *Phys.Rev.* **D61** 095009 (2000); *hep-ph/9911365*.
- [7] D.Bailey, B.Campbell, S.Davidson, *Z.Phys.* **C61** 613 (1994); *hep-ph/9309310*.
- [8] J.Ellis, G.Leontaris, J.Rizos, *JHEP* 0005 (2000) 001; *hep-ph/0002263*.
- [9] R.Barbieri, A.Strumia, Z.Berezhiani, *Phys.Lett.* **B407** 250 (1997); *hep-ph/9704275*.
- [10] G.Bhattacharyya, A.Raychaudhuri, *Phys.Rev.* **D57** 3837 (1998); *hep-ph/9712245*.
- [11] J.Jang, J.Kim, J.Lee, *Phys.Rev.* **D55** 7296 (1997); *hep-ph/9701283*.
- [12] S.Baek, Y.Kim, *Phys.Rev.* **D60** 077701 (1999); *hep-ph/9906385*.
- [13] W.Marciano, A.Sirlin, *Phys.Rev.Lett.* **71** 3629 (1993).
- [14] M.Finkemeier, *Phys.Lett.* **B387** 391 (1996); *hep-ph/9505434*.
- [15] D.Groom et al. (Particle Data Group), *Eur.Phys.J.* **C15** 1 (2000).
We have taken into account the 2001 update from Dec.12.
- [16] F.Ledroit, G.Sajot, *ISN Grenoble*, **GDR-S-008**.
- [17] T.Bowder et al. (CLEO Collaboration), *Int.J.Mod.Phys.A* **1651B** 636 (2001); *hep-ex/0007057*.
- [18] H.Dreiner, P.Morawitz, *Nucl.Phys.* **B503** 55 (1997); *hep-ph/9703279*.

[19] A.Buras, *hep-ph/9806471*.

[20] M.Neubert, *hep-ph/0110301*.

[21] C.Bernard, *Nucl.Phys.(Proc.Suppl.)* **B94** 159 (2001); *hep-lat/0011064*.

lower limit $(m_{\text{SUSY}}/100 \text{ GeV})^2$	product of $\mathcal{R}_p$ coupling constants	upper limit $(m_{\text{SUSY}}/100 \text{ GeV})^2$	exchanged sfermion
-7.9×10^{-8}	$\text{Re}[\lambda'_{k11} \lambda_{1k1}]$	7.1×10^{-5}	$\widetilde{\ell}_L^k$
-7.9×10^{-5}	$\text{Re}[\lambda'_{311} \lambda_{232}]$	—	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k11} \lambda_{3k1} $	3.4×10^{-6}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{211} \lambda_{322} $	1.5×10^{-3}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{111} \lambda_{211} $	3.4×10^{-6}	$\widetilde{\ell}_L^k$
-7.0×10^{-7}	$\text{Re}[\lambda'_{k12} \lambda_{1k1}]$	1.8×10^{-5}	$\widetilde{\ell}_L^k$
-1.8×10^{-4}	$\text{Re}[\lambda'_{k12} \lambda_{2k2}]$	$3.8 \times 10^{-3}, k = 3$	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k12} \lambda_{2k1} $	5.4×10^{-6}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k12} \lambda_{3k1} $	5.4×10^{-6}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k12} \lambda_{1k2} $	1.3×10^{-3}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k12} \lambda_{3k2} $	1.3×10^{-3}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{313} \lambda_{233} $	2×10^{-3}	$\widetilde{\ell}_L^k$
-6.4×10^{-4}	$\text{Re}[\lambda'_{213} \lambda_{323}]$	1×10^{-3}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k13} \lambda_{3k1} $	6×10^{-4}	$\widetilde{\ell}_L^k$
0	$ \lambda'_{k13} \lambda_{3k2} $	7×10^{-4}	$\widetilde{\ell}_L^k$

$\frac{\text{lower limit}}{(m_{\text{Susy}}/100 \text{ GeV})^2}$	product of $\mathcal{R}_p$ coupling constants	$\frac{\text{upper limit}}{(m_{\text{Susy}}/100 \text{ GeV})^2}$	exchanged sfermion
0	$ \lambda'_{123} \lambda_{222}^* $	8×10^{-3}	$\widetilde{u_L^k}$
0	$ \lambda'_{1k2} \lambda_{2k3}^* , k \neq 1$	8×10^{-3}	$\widetilde{u_L^k}$
0	$ \lambda'_{1k2} \lambda_{2k1}^* $	3×10^{-7}	$\widetilde{u_L^k}$
0	$ \lambda'_{1k1} \lambda_{2k2}^* $	3×10^{-7}	$\widetilde{u_L^k}$
0	$ \lambda_{k32}^* \lambda_{k12} $	7×10^{-5}	$\widetilde{\nu_L^k}$
0	$ \lambda_{k23}^* \lambda_{k21} $	7×10^{-5}	$\widetilde{\nu_L^k}$
0	$ \lambda_{k32}^* \lambda_{k21} $	7×10^{-5}	$\widetilde{\nu_L^k}$
0	$ \lambda_{k23}^* \lambda_{k12} $	7×10^{-5}	$\widetilde{\nu_L^k}$
0	$ \lambda_{k21}^* \lambda_{k12} $	6×10^{-9}	$\widetilde{\nu_L^k}$
0	$ \lambda_{k12}^* \lambda_{k12} $	6×10^{-9}	$\widetilde{\nu_L^k}$
0	$ \lambda'_{k12} \lambda_{k21}^* $	6×10^{-9}	$\widetilde{\nu_L^k}$
0	$ \lambda'_{k21} \lambda_{k21}^* $	6×10^{-9}	$\widetilde{\nu_L^k}$
0	$ \lambda_{311}^* \lambda_{312} $	3×10^{-3}	$\widetilde{\nu_L^k}$
0	$ \lambda'_{311} \lambda_{321}^* $	3×10^{-3}	$\widetilde{\nu_L^k}$